

Table XVII-7
MESOTHELIOMASURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Relative Survival Rates (SEER ^b)																						
1-year	42.5	43.9	44.1	53.3	35.1	47.0	38.8	53.1	50.7	60.3	25.9	57.2	39.2	32.6	48.6	55.9	41.9	39.7	54.6	36.1	50.6	42.0
2-year	31.5	29.9	21.2	38.7	18.2	25.5	19.9	27.2	33.1	41.8	14.5	44.3	23.6	27.3	27.8	39.2	37.5	19.2	35.6	20.0	32.0	
3-year	24.8	20.5	15.6	27.8	10.8	23.0	17.3	20.9	24.6	34.1	12.3	35.3	21.7	22.2	17.4	26.3	35.8	17.1	24.0	10.3		
4-year	22.2	15.0	9.7	26.0	10.8	20.9	8.9	11.9	24.6	30.3	10.3	33.9	20.1	14.9	14.7	22.1	34.0	8.8	20.3			
5-year	20.9	13.9	9.7	19.7	8.6	20.9	8.9	11.9	23.6	30.3	10.3	29.6	18.5	11.0	14.7	22.1	26.8	0.0				
6-year	18.1	11.3	9.7	15.3	8.6	18.9	8.9	7.5	19.0	26.2	10.3	28.0	16.2	9.2	14.7	22.1	21.4					
7-year	17.4	11.3	9.7	12.9	8.6	16.3	8.9	7.5	16.7	26.2	10.3	28.0	16.2	7.1	10.1	21.6						
8-year	17.1	11.3	7.5	10.8	8.6	13.4	6.8	7.5	14.4	26.2	10.3	26.5	16.2	7.1	10.1							
9-year	17.1	10.4	7.5	10.8	8.6	13.4	6.8	7.5	12.6	25.4	10.3	26.5	15.8	5.2								
11-year	16.2	8.6	4.1	10.8	8.6	13.4	6.8	7.5	12.6	22.8	10.3	26.3	15.8									
10-year	15.7	8.6	4.1	10.8	8.6	13.4	6.8	7.5	12.6	19.8	10.3	26.3										
12-year	15.7	8.6	0.0	10.8	8.6	13.4	6.8	7.5	12.6	16.0	10.3											
13-year	15.7	8.6	0.0	10.8	8.6	13.4	6.8	7.5	12.6	16.0												
14-year	15.7	8.6	0.0	10.8	8.6	13.4	6.8	7.5	12.6													
15-year	15.7	8.6	0.0	10.8	8.6	13.4	6.8	7.5														
16-year	15.7	8.6	0.0	10.8	8.6	13.4	6.8															
17-year	15.6	8.6	0.0	10.8	8.6	13.4																
18-year	15.6	8.6	0.0	10.8	8.6																	
19-year	15.6	8.6	0.0	10.8																		
20-year	15.6	8.6	0.0																			

^a

Survival rates are relative rates expressed as percents.

^b

Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).