

Table XVII-6

MESOTHELIOMASURVIVAL RATES^a

By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Relative Survival Rates (SEER ^b)																						
1-year	35.8	37.5	41.6	35.3	40.4	35.9	29.8	37.2	31.4	36.6	35.4	34.7	36.3	46.3	34.2	35.1	32.0	42.0	39.1	31.8	46.6	38.1
2-year	12.0	16.6	21.1	11.4	13.5	13.6	14.9	12.8	8.6	15.4	14.0	11.5	11.2	25.9	14.6	18.4	14.6	14.3	21.7	16.3	18.1	
3-year	6.7	9.8	14.0	6.4	7.9	8.3	7.8	7.8	8.0	8.1	8.1	7.0	9.2	16.3	9.0	12.7	7.7	8.9	12.2	6.1		
4-year	5.7	7.1	10.0	3.8	4.6	7.0	6.1	4.5	4.3	6.4	5.3	4.0	6.1	11.3	6.6	10.5	5.4	6.5	9.6			
5-year	4.2	5.6	6.6	3.8	3.7	6.4	4.2	4.5	4.3	4.8	2.4	3.5	4.4	10.4	4.5	10.5	4.8	3.8				
6-year	3.5	4.7	5.0	3.1	3.0	2.9	3.0	3.3	4.3	2.9	2.4	3.5	3.7	10.4	2.4	8.6	4.8					
7-year	3.1	4.0	3.1	3.1	2.1	2.0	3.0	1.7	4.3	2.3	2.4	3.0	2.3	9.8	1.7	8.6						
8-year	2.5	3.8	3.1	3.1	2.1	2.0	3.0	0.9	4.3	2.3	1.0	2.4	2.3	7.1	1.7							
9-year	2.3	3.8	2.3	3.1	1.2	2.0	3.0	0.9	2.9	2.3	1.0	2.4	0.9	5.6								
10-year	2.3	3.8	2.3	2.7	1.2	2.0	2.8	0.9	2.9	1.9	1.0	2.4	0.0									
11-year	2.3	3.8	2.3	2.7	1.2	2.0	2.8	0.9	2.2	1.9	1.0	2.4										
12-year	2.3	3.8	2.3	2.7	1.2	2.0	2.8	0.9	2.2	1.9	1.0											
13-year	2.0	3.8	2.3	2.7	1.2	1.4	2.8	0.9	2.2	1.9												
14-year	2.0	3.8	2.3	2.7	1.2	0.0	2.8	0.9	2.2													
15-year	1.4	3.8	2.3	2.7	0.0	0.0	2.8	0.9														
16-year	1.4	3.8	2.3	2.7	0.0	0.0	2.8															
17-year	1.4	3.8	2.3	2.7	0.0	0.0																
18-year	1.4	3.8	2.3	2.7	0.0																	
19-year	1.4	3.8	2.3	2.7																		
20-year	1.4	3.8	2.3																			

^a Survival rates are relative rates expressed as percents.^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).